1 The Swendsen–Wang method and its generalizations

Several enhancements of Monte Carlo methods are based on a remarkable trick: take a big and difficult problem, and replace it by an even bigger problem that contains the first problem; then solve the bigger problem with methods that were not applicable to the first.

The Swendsen–Wang method takes a spin system such as

$$P(\mathbf{x}) = \frac{1}{Z} \exp\left(\sum_{n,n'} J_{nn'} x_n x_{n'}\right),\tag{1}$$

and replaces it by a bigger system that contains both the N original spin variables and M additional 'bond' variables, where M is the number of non-zero couplings $J_{nn'}$ in the original spin system.

The extended Ising model

Let's represent the original Ising model by the product of factors

$$P(\mathbf{x}) = \frac{1}{Z} \prod_{m=1}^{M} f_m(\mathbf{x}_m), \qquad (2)$$

where each factor $f_m(\mathbf{x}_m)$ depends on just two of the spins, say x_n and $x_{n'}$:

$$f_m(\mathbf{x}_m) = f(x_n, x_{n'}) = \exp(J_{nn'} x_n x_{n'}).$$
 (3)

In a standard Ising model, all the couplings between neighbouring spins $J_{nn'}$ are equal to a single positive number, J. Later, we will generalize our description to arbitrary spin systems.

We extend this model by introducing M additional 'bond' variables, **d**.



spins

spins and bonds

We define a joint distribution that couples the spins to the bonds,

$$P(\mathbf{x}, \mathbf{d}) = \frac{1}{Z'} \prod_{m=1}^{M} g_m(\mathbf{x}_m, d_m), \qquad (4)$$

in such a way that

- (a) The marginal distribution of \mathbf{x} , $\sum_{\mathbf{d}} P(\mathbf{x}, \mathbf{d})$, is equal to the Ising distribution (2);
- (b) the conditional distributions $P(\mathbf{x}|\mathbf{d})$ and $P(\mathbf{d}|\mathbf{x})$ are both simple to sample from.

Each bond variable has two possible states: 1 (also known as 'open' or 'connected') and 0 ('closed' or 'disconnected'). The factor $g_m(\mathbf{x}_m, d_m)$ is defined by the following

eight values:

$$g_{m}(\mathbf{x}_{m}, d_{m}) = \begin{cases} d_{m} = 0 & d_{m} = 1 \\ x_{n'} = 0 & x_{n'} = 1 & x_{n'} = 0 & x_{n'} = 1 \\ x_{n} = 0 & e^{-J} & e^{-J} & e^{J} - e^{-J} & 0 \\ x_{n} = 1 & e^{-J} & e^{-J} & 0 & e^{J} - e^{-J}, \end{cases}$$
(5)

The distribution of \mathbf{x} and \mathbf{d} is unchanged if we rescale any factor by any constant; it will be convenient to introduce an alternative notation for the factor g_m , rescaling it by e^J , and defining

$$p \equiv 1 - e^{-2J}.\tag{6}$$

The rescaled factor is:

$$\tilde{g}_{m}(\mathbf{x}_{m}, d_{m}) = \begin{cases}
 d_{m} = 0 & d_{m} = 1 \\
 x_{n'} = 0 & x_{n'} = 1 & x_{n'} = 0 & x_{n'} = 1 \\
 x_{n} = 0 & 1 - p & 1 - p & p & 0 \\
 x_{n} = 1 & 1 - p & 1 - p & 0 & p,
 \end{cases}$$
(7)

The extended model can thus be written as

$$P(\mathbf{x}, \mathbf{d}) = \frac{1}{Z} \prod_{m} g_m(\mathbf{x}_m, d_m) = \frac{1}{\tilde{Z}} \prod_{m} \tilde{g}_m(\mathbf{x}_m, d_m).$$
(8)

The first partition function Z is identical to the partition function of the original Ising model. The second is

$$\tilde{Z} = Z e^{MJ} \tag{9}$$

(Check sign.)

In the case where the original spins are connected in a rectangular grid, the factor graph for the extended model looks like this:



1.1 Sampling from the extended model by Gibbs sampling

The conditional distributions $P(\mathbf{x}|\mathbf{d})$ and $P(\mathbf{d}|\mathbf{x})$ are as follows:

 $P(\mathbf{x}|\mathbf{d})$ – the bonds connect the spins into a number of clusters (connected components); all spins in a cluster must adopt the same state as each other; the states ± 1 are selected with equal probability.

 $P(\mathbf{d}|\mathbf{x})$ – conditional on the spins, the bonds are independent. If the two spins surrounding a bond are equal $(x_n = x_{n'})$, set the bond d_m to 1 with probability p (see equation (6)); otherwise set it to zero.

In the case of a rectangular grid, this Gibbs sampling algorithm mixes very rapidly.

The following figures illustrate Gibbs sampling. Spin states up and down are shown by filled and empty circles. Bond states 1 and 0 are shown by thick lines and thin dotted lines. We start from a state with five connected components. (Remember that isolated spins count as connected components, albeit of size 1.)



First, let's update the bonds The forbidden bonds are highlighted Bonds are forbidden from forming wherever the two adjacent spins are in opposite



1.2 Other properties of the extended model

We already mentioned that the partition function Z is the same as that of the Ising model.

The marginal $P(\mathbf{x})$ is correct, because when we sum the factor g_m over d_m , we get f_m . Summing over d_m is easy because it appears in only one factor.

OK, we've summed out **d** and obtained the Ising model. What if we sum out **x**? The marginal $P(\mathbf{d})$ is called the *random cluster model*. Summing over **x** for given **d**, all factors are constants. The number of states is 2^{number of clusters}. Thus

$$P(\mathbf{d}) = \frac{1}{\tilde{Z}} \prod_{m} \left(p^{d_m} (1-p)^{1-d_m} \right) 2^{c(\mathbf{d})}$$
(10)

where $c(\mathbf{d})$ is the number of connected components in the state \mathbf{d} . Isolated spins whose neighbouring bonds are all zero count as single connected components.

The random cluster model can be generalized by replacing the number 2 by a parameter q:

$$P^{(q)}(\mathbf{d}) = \prod_{m} \left(p^{d_m} (1-p)^{1-d_m} \right) q^{c(\mathbf{d})}$$
(11)

The random cluster model can be simulated directly, just as the Ising model can be simulated directly; but the S–W method, augmenting the bonds with spins, is probably the most efficient way to simulate the model. For integer values of q, the appropriate spin system is the 'Potts model', the generalization of the Ising model from 2 spin states to q.

2 S–W for General spin system

We now include a bias h_n at each spin. And we allow the couplings J to be positive or negative.

Assuming that the original coupling associated with bond m, $J_{nn'}$, is positive, the factor $g_m(\mathbf{x}_m, d_m)$ is defined by the following eight values:

$$g_m(\mathbf{x}_m, d_m) = \begin{cases} d_m = 0 & d_m = 1 \\ x_{n'} = 0 & x_{n'} = 1 & x_{n'} = 0 & x_{n'} = 1 \\ x_n = 0 & e^{-J_{nn'}} & e^{-J_{nn'}} & e^{J_{nn'}} - e^{-J_{nn'}} & 0 \\ x_n = 1 & e^{-J_{nn'}} & e^{-J_{nn'}} & 0 & e^{J_{nn'}} - e^{-J_{nn'}}, \end{cases}$$
(12)

As before we can introduce a parameter $p_m \equiv 1 - e^{-2J_{nn'}}$ and write a rescaled factor:

$$\tilde{g}_{m}(\mathbf{x}_{m}, d_{m}) = \begin{cases}
d_{m} = 0 & d_{m} = 1 \\
x_{n'} = 0 & x_{n'} = 1 & x_{n'} = 0 & x_{n'} = 1 \\
x_{n} = 0 & 1 - p & 1 - p & p & 0 \\
x_{n} = 1 & 1 - p & 1 - p & 0 & p,
\end{cases}$$
(13)

If $J_{nn'}$ is *negative*, we define the factor g_m thus:

$$g_{m}(\mathbf{x}_{m}, d_{m}) = \begin{cases} d_{m} = 0 & d_{m} = 1 \\ x_{n'} = 0 & x_{n'} = 1 & x_{n'} = 0 & x_{n'} = 1 \\ x_{n} = 0 & e^{J_{nn'}} & e^{J_{nn'}} & 0 & e^{-J_{nn'}} - e^{J_{nn'}} \\ x_{n} = 1 & e^{J_{nn'}} & e^{J_{nn'}} & e^{-J_{nn'}} - e^{J_{nn'}} & 0, \end{cases}$$
(14)

Two spins surrounding such a bond must be in *opposite* states if the bond is connected $(d_m = 1)$.